

Solving large classes of nonlinear systems of PDEs

Roumen Anguelov, Elemér E Rosinger

Department of Mathematics
and Applied Mathematics
University of Pretoria
Pretoria
0002 South Africa
eerosinger@hotmail.com

Abstract

It is shown that large classes of nonlinear systems of PDEs, with possibly associated initial and/or boundary value problems, can be solved by the method of order completion. The solutions obtained can be assimilated with Hausdorff continuous functions. The usual Navier-Stokes equations, as well as their various modifications aiming at a realistic modelling, are included as particular cases. The same holds for the critically important constitutive relations in various branches of Continuum Mechanics. The solution method does not involve functional analysis, nor various Sobolev or other spaces of distributions or generalized functions. The general and type independent *existence* and *regularity* results regarding solutions presented here have recently been introduced in the literature.

”... provided also if need be that the notion of a solution
shall be suitably extended ...”

cited from Hilbert’s 20th Problem

1. Main ideas of the order completion solution method

The solution method is divided in two parts. The proof of the *existence*

of solutions follows the method of order completion introduced and first developed in Oberguggenberger & Rosinger. The proof of the *regularity* of solutions is a consequence of recent results obtained in Anguelov [1], regarding the structure of the Dedekind order completion of spaces of continuous functions $\mathcal{C}(X)$, where X is a rather arbitrary topological space.

For simplicity of presentation, we shall consider single nonlinear PDEs. The extension to systems of such nonlinear PDEs and associated initial and/or boundary value problems can - rather surprisingly - be done easily, as seen in Oberguggenberger & Rosinger, this is being one of the major advantages of the order completion method. Let us therefore consider nonlinear PDEs of the general form

$$(1.1) \quad F(x, U(x), \dots, D_x^p U(x), \dots) = f(x), \quad x \in \Omega \subseteq \mathbb{R}^n$$

with $p \in \mathbb{N}$, $|p| \leq m$, where the domains Ω can be any open, not necessarily bounded subsets of $\mathbb{R}^n$, while the orders $m \in \mathbb{N}$ of the PDEs are arbitrary given, and the unknown functions, that is, the solutions one looks for are $U : \Omega \longrightarrow \mathbb{R}$.

The *unprecedented generality* of these nonlinear PDEs comes, above all, from the class of functions F which define the left hand terms, and which are only assumed to be *jointly continuous* in all of their arguments. The right hand terms f are also required to be *continuous*.

As seen, however, both F and f can have certain *discontinuities* as well, Oberguggenberger & Rosinger.

Regardless of the above generality of the nonlinear systems of PDEs considered, one can find for them solutions U defined on the *whole* of the respective domains Ω . These solutions U have the *blanket, type independent*, or *universal regularity* property that they can be assimilated with *Hausdorff continuous functions*.

It follows in this way that, when solving systems of nonlinear PDEs of the generality of those in (1.1), one can *dispense with* the various customary spaces of distributions, hyperfunctions, generalized functions, Sobolev spaces, and so on. Instead one can stay within the realms of *usual functions*, more precisely, of *interval valued* functions, see the Appendix for a short presentation of essentials on Hausdorff continuous functions. Also, when proving the *existence* and the mentioned type of *regularity* of such solutions one can dispense with methods of Functional Analysis. However, functional analytic methods can pos-

sibly be used in order to obtain further regularity or other desirable properties of such solutions.

Let us now associate with each nonlinear PDE in (1.1) the corresponding nonlinear partial differential operator defined by the left hand side, namely

$$(1.2) \quad T(x, D)U(x) = F(x, U(x), \dots, D_x^p U(x), \dots), \quad x \in \Omega$$

Two facts about the nonlinear PDEs in (1.1) and the corresponding nonlinear partial differential operators $T(x, D)$ in (1.2) are important and immediate :

- The operators $T(x, D)$ can *naturally* be seen as acting in the *classical* context, namely

$$(1.3) \quad T(x, D) : \mathcal{C}^m(\Omega) \ni U \longmapsto T(x, D)U \in \mathcal{C}^0(\Omega)$$

while, unfortunately on the other hand :

- The mappings in this natural classical context (1.3) are typically *not* surjective even in the case of linear $T(x, D)$, and they are even less so in the general nonlinear case of (1.1), (1.2).

In other words, linear or nonlinear PDEs in (1.1) typically *cannot* be expected to have *classical* solutions $U \in \mathcal{C}^m(\Omega)$, for arbitrary continuous right hand terms $f \in \mathcal{C}^0(\Omega)$, as illustrated by a variety of well known examples, some of them rather simple ones, see Oberguggenberger & Rosinger [chap. 6].

Furthermore, it can often happen that nonclassical solutions do have a major applicative interest, thus they have to be sought out *beyond* the confines of the classical framework in (1.3).

This is, therefore, how we are led to the *necessity* to consider *generalized solutions* U for PDEs like those in (1.1), that is, solutions $U \notin \mathcal{C}^m(\Omega)$, which therefore are no longer classical. This means that the natural classical mappings (1.3) must in certain suitable ways be *extended* to *commutative diagrams*

$$(1.4) \quad \begin{array}{ccc} \mathcal{C}^m(\Omega) & \xrightarrow{T(x, D)} & \mathcal{C}^0(\Omega) \\ \downarrow \subseteq & & \downarrow \subseteq \\ X & \xrightarrow{\tilde{T}} & Y \end{array}$$

with the generalized solutions now being found as

$$(1.5) \quad U \in X \setminus \mathcal{C}^m(\Omega)$$

instead of the classical ones $U \in \mathcal{C}^m(\Omega)$ which may easily fail to exist. A further important point is that one expects to reestablish certain kind of *surjectivity* type properties typically missing in (1.3), at least such as for instance

$$(1.6) \quad \mathcal{C}^0(\Omega) \subseteq \tilde{T}(X)$$

As it turns out, when constructing extensions of (1.3) given by commutative diagrams (1.4), we shall be interested in the following somewhat larger spaces of piecewise smooth functions. For any integer $0 \leq l \leq \infty$, we define

$$(1.7) \quad \mathcal{C}_{nd}^l(\Omega) = \left\{ u : \Omega \rightarrow \mathbb{R} \left| \begin{array}{l} \exists \Gamma \subset \Omega \text{ closed, nowhere dense} : \\ u \in \mathcal{C}^l(\Omega \setminus \Gamma) \end{array} \right. \right\}$$

and as an immediate strengthening of (1.3), we obviously obtain

$$(1.8) \quad T(x, D) \mathcal{C}_{nd}^m(\Omega) \subseteq \mathcal{C}_{nd}^0(\Omega)$$

The solution of the nonlinear PDEs in (1.1) through the order completion method will come from the construction of specific instances of the *commutative diagrams* (1.4), given by, see (2.18), (2.27)

$$\begin{array}{ccc}
\mathcal{C}_{nd}^m(\Omega) & \xrightarrow{T(x, D)} & \mathcal{C}_{nd}^0(\Omega) \\
\downarrow & & \downarrow \\
(1.9) \quad \mathcal{M}_T^m(\Omega) & \xrightarrow{T} & \mathcal{M}^0(\Omega) \\
\downarrow & & \downarrow \\
\mathcal{M}_T^m(\Omega)^\# & \xrightarrow[\text{bijective}]{T^\#} & \mathcal{M}^0(\Omega)^\#
\end{array}$$

where, as elaborated later, the operation $(\)^\#$ means the *Dedekind order completion*, according to MacNeille, of the respective spaces, as well as the extension to such completions of the respective mappings, see Oberguggenberger & Rosinger [Appendix]. It follows that in terms of (1.4), we have

$$X = \mathcal{M}_T^m(\Omega)^\#, \quad Y = \mathcal{M}^0(\Omega)^\#, \quad \tilde{T} = T^\#$$

thus we shall obtain for the nonlinear PDEs in (1.1) generalized solutions

$$(1.10) \quad U \in \mathcal{M}_T^m(\Omega)^\#$$

Furthermore, instead of the *surjectivity* condition (1.6), we shall at least have the following stronger one

$$(1.11) \quad \mathcal{C}_{nd}^0(\Omega) \subseteq T^\#(\mathcal{M}_T^m(\Omega)^\#)$$

So far about the main ideas related to the *existence* of solutions of general nonlinear PDEs of the form (1.1).

As for the *regularity* of such solutions, we recall that, as shown in Oberguggenberger & Rosinger, one has the inclusions

$$(1.12) \quad \mathcal{M}^0(\Omega)^\# \subseteq Mes(\Omega)$$

where $Mes(\Omega)$ denotes the set of Lebesgue measurable functions on Ω . In this way, in view of (1.9), (1.10), one can assimilate the generalized solutions U of the nonlinear PDEs in (1.1) with usual measurable functions in $Mes(\Omega)$.

Recently, however, based on results in Anguelov [1], it was shown that instead of (1.12), one has the much stronger property

$$(1.13) \quad \mathcal{M}^0(\Omega)^\# \subseteq \mathbb{H}(\Omega)$$

where $\mathbb{H}(\Omega)$ denotes the set of Hausdorff continuous functions on Ω . Consequently, now one can significantly improve on the earlier regularity result, as one can assimilate the generalized solutions U of the nonlinear PDEs in (1.1) with usual functions in $\mathbb{H}(\Omega)$.

2. The construction of diagram (1.9)

Since we solve PDEs through order completion, let us see how near we can come to satisfying the equality in (1.1), when using inequalities. For that purpose, it is useful to consider for each $x \in \Omega$ the following set of real numbers

$$(2.1) \quad \mathbb{R}_x = \{ F(x, \xi_0, \dots, \xi_p, \dots) \mid \xi_p \in \mathbb{R}, \text{ for } p \in \mathbb{N}^n, |p| \leq m \}$$

Clearly, for $x \in \Omega$ fixed, $\mathbb{R}_x$ is the range in $\mathbb{R}$ of $F(x, \dots)$, and since F is jointly continuous in all its arguments, it follows that $\mathbb{R}_x$ is a nonvoid interval which is bounded, half bounded, or is the whole of $\mathbb{R}$. This latter case, which can happen often with nonlinear PDEs in (1.1), will be easier to deal with, see (2.3) next. Clearly, in the case of non-degenerate linear PDEs in (1.1), this latter case always happens. Given now $x \in \Omega$, it is obvious that a *necessary* condition for the existence of a classical smooth solution $U \in \mathcal{C}^m$ of (1.1) in a neighbourhood of x is the condition

$$(2.2) \quad f(x) \in \mathbb{R}_x$$

Consequently, for the time being, we shall make the assumption that the right hand term functions f in the nonlinear PDEs in (1.1) satisfy the somewhat stronger version of condition (2.2) given by

$$(2.3) \quad f(x) \in \text{int } \mathbb{R}_x, \text{ for } x \in \Omega$$

Clearly, whenever we have

$$(2.4) \quad \mathbb{R}_x = \mathbb{R}, \text{ for } x \in \Omega$$

then (2.3) is satisfied. And as mentioned, this is the case with all nontrivial linear PDEs, as well as with most of the nonlinear PDEs of practical interest.

And now the basic and rather simple *local approximation* result on how near we can satisfy the equality in (1.1), when using inequalities.

Proposition 2.1.

Given $f \in \mathcal{C}^0(\Omega)$, then

$$(2.5) \quad \begin{aligned} & \forall x_0 \in \Omega, \quad \epsilon > 0 : \\ & \exists \delta > 0, \quad P \text{ polynomial in } x \in \mathbb{R}^n : \\ & \forall x \in \Omega, \quad \|x - x_0\| \leq \delta : \\ & \quad f(x) - \epsilon \leq T(x, D)P(x) \leq f(x) \end{aligned}$$

Proof

Given $x_0 \in \Omega$, then for $\epsilon > 0$ small enough, condition (2.3) yields $\xi_p \in \mathbb{R}$, with $p \in \mathbb{N}^n$, $|p| \leq m$, such that

$$(2.6) \quad F(x_0, \xi_0, \dots, \xi_p, \dots) = f(x_0) - \epsilon/2$$

Let us take P a polynomial in $x \in \mathbb{R}^n$, which satisfies the conditions

$$D_x^p P(x_0) = \xi_p, \quad p \in \mathbb{N}^n, \quad |p| \leq m$$

In this case from (2.6) we clearly obtain the relation

$$(2.7) \quad T(x_0, D)P(x_0) = f(x_0) - \epsilon/2$$

and since both $T(x, D)P(x)$ and $f(x)$ are continuous in $x \in \Omega$, the local inequality property (2.5) follows easily from (2.7).

The *global approximation* version of the inequality property in (2.5) is given in

Proposition 2.2.

If $f \in \mathcal{C}^0(\Omega)$, then

$$(2.8) \quad \begin{aligned} & \forall \epsilon > 0 : \\ & \exists \Gamma_\epsilon \subset \Omega \text{ closed, nowhere dense, } U_\epsilon \in \mathcal{C}^m(\Omega \setminus \Gamma_\epsilon) : \\ & \quad f - \epsilon \leq T(x, D)U_\epsilon \leq f \text{ on } \Omega \setminus \Gamma_\epsilon \end{aligned}$$

Proof

Let us take a covering of Ω of the form

$$(2.9) \quad \Omega = \bigcup_{\nu \in \mathbb{N}} K_\nu$$

where K_ν are compact n-dimensional intervals in $\mathbb{R}^n$, namely, $K_\nu = [a_\nu, b_\nu]$, with $a_\nu = (a_{\nu,1}, \dots, a_{\nu,n})$, $b_\nu = (b_{\nu,1}, \dots, b_{\nu,n}) \in \mathbb{R}^n$. We also assume, see Forster, that the covering (2.9) is locally finite, that is

$$(2.10) \quad \begin{aligned} & \forall x \in \Omega : \\ & \exists V_x \subseteq \Omega \text{ neighbourhood of } x : \\ & \{ \nu \in \mathbb{N} \mid K_\nu \cap V_x \neq \emptyset \} \text{ is a finite set of indices} \end{aligned}$$

and furthermore

$$(2.11) \quad \text{the interiors of } K_\nu, \text{ with } \nu \in \mathbb{N}, \text{ are pairwise disjoint}$$

Let us now take $\epsilon > 0$ arbitrary but fixed. Further, we take $\nu \in \mathbb{N}$. We shall apply Proposition 2.1 to each $x_0 \in K_\nu$. Then we obtain $\delta_{x_0} > 0$ and a polynomial P_{x_0} such that

$$f(x) - \epsilon \leq T(x, D)P_{x_0}(x) \leq f(x), \quad x \in \Omega, \quad \|x - x_0\| \leq \delta_{x_0}$$

But K_ν is compact, therefore

$$(2.12) \quad \begin{aligned} & \exists \delta > 0 : \\ & \forall x_0 \in K_\nu : \\ & \exists P_{x_0} \text{ polynomial in } x \in \mathbb{R}^n : \\ & \forall x \in \Omega, \quad \|x - x_0\| \leq \delta_{x_0} : \\ & f(x) - \epsilon \leq T(x, D)P_{x_0}(x) \leq f(x) \end{aligned}$$

Now we shall subdivide K_ν , which was assumed to be a compact n-dimensional interval, into n-dimensional subintervals $I_1, \dots, I_\mu$, so that the diameter of each of them is less or equal δ .

Let us denote by J a generic such n-dimensional subinterval in any of the K_ν , when $\nu \in \mathbb{N}$. If $a_J \in J$ is the center of any such n-dimensional subinterval then (2.12) gives for $x \in \text{int } J$

$$f(x) - \epsilon \leq T(x, D)P_{a_J}(x) \leq f(x)$$

Let us now take

$$(2.13) \quad \Gamma_\epsilon = \Omega \setminus \bigcup_J \text{int } J$$

that is, with the union ranging over all such n -dimensional subintervals J . In view of ??? If we define $U_\epsilon \in \mathcal{C}^m(\Omega \setminus \Gamma_\epsilon)$ by

$$U_\epsilon = P_{a,J} \quad \text{on } \Omega \cap \text{int } J$$

then the proof is completed.

Remark 2.1

1) It is easy to see that the inequalities in (2.5) and (2.8) can be replaced with the following ones, respectively

$$(2.14) \quad f(x) \leq T(x, D)P(x) \leq f(x) + \epsilon$$

$$(2.15) \quad f \leq T(x, D)U_\epsilon \leq f + \epsilon$$

as their proofs follow after the corresponding obvious minor changes in the proofs of the above two propositions. And these four inequalities are *sharper* than would respectively be the inequalities

$$f(x) - \epsilon \leq T(x, D)P(x) \leq f(x) + \epsilon, \quad f - \epsilon \leq T(x, D)U_\epsilon \leq f + \epsilon$$

As we shall see not much later, we do need the sharper inequalities. Indeed, the order completion method which we shall employ is based on MacNeille's construction, therefore, it uses *Dedekind cuts*. And such cuts do need the above sharper inequalities.

2) In Proposition 2.2., as well as in its version corresponding to the above inequality (2.15), we have in addition the property

$$(2.16) \quad \text{mes}(\Gamma_\epsilon) = 0$$

where *mes* denotes the usual Lebesgue measure. Indeed, according to (2.10), (2.11) and (2.13), Γ_ϵ is a countable union of rectangular grids, each generated by a finite number of hyperplanes.

Here it should be noted that the presence of the *closed, nowhere dense* singularity sets Γ_ϵ in the *global* inequalities (2.8) and (2.15) will prove not to be a hindrance. And in fact, it will lead to the classes of piecewise smooth functions in (1.7) which prove to be convenient.

The presence of such closed, nowhere dense *singularity* sets is rather deep rooted, as it is connected with such facts as the *flabbiness* of related sheaves of functions, see Oberguggenberger & Rosinger [chapter 7], or the global version of the classical Cauchy-Kovalevskia theorem

on analytic nonlinear PDEs, see Oberguggenberger & Rosinger and the literature cited there.

3) As seen from the proof of Proposition 2.2., the functions U_ϵ can in fact be chosen as piecewise polynomials in $x \in \mathbb{R}^n$.

Let us now note that there is an obvious ambiguity with the piecewise smooth functions in $\mathcal{C}_{nd}^l(\Omega)$ in (1.7). Indeed, given any such function $u \in \mathcal{C}_{nd}^l(\Omega)$, the corresponding closed, nowhere dense set Γ cannot be defined uniquely. Therefore, it is convenient to factor out this ambiguity, and this can be done easily as follows. Since $\mathcal{C}_{nd}^0(\Omega)$ is the largest of these spaces of functions, we shall do for this space the mentioned factoring out, by defining on it the *equivalence* relation $u \approx v$ for any two elements $u, v \in \mathcal{C}_{nd}^0(\Omega)$, as given by the condition

$$(2.17) \quad \begin{aligned} & \exists \Gamma \subset \Omega \text{ closed, nowhere dense} : \\ & *) \quad u, v \in \mathcal{C}^0(\Omega \setminus \Gamma) \\ & **) \quad u = v \text{ on } \Omega \setminus \Gamma \end{aligned}$$

It is easy to see that $\approx$ defined above is indeed an equivalence relation, since the union of a finite number of closed and nowhere dense subsets is again closed and nowhere dense. Now we can eliminate the mentioned ambiguity by going to the *quotient* space

$$(2.18) \quad \mathcal{M}^0(\Omega) = \mathcal{C}_{nd}^0(\Omega) / \approx$$

and in view of (2.8), (2.15), we define for any two elements $G, H \in \mathcal{M}^0(\Omega)$ the partial order $G \leq H$, by

$$(2.19) \quad \begin{aligned} & \exists g \in G, h \in H, \Gamma \subset \Omega \text{ closed, nowhere dense} : \\ & *) \quad g, h \in \mathcal{C}^0(\Omega \setminus \Gamma) \\ & **) \quad g \leq h \text{ on } \Omega \setminus \Gamma \end{aligned}$$

Let us now denote by

$$(2.20) \quad (\mathcal{M}^0(\Omega)^\#, \leq)$$

the Dedekind order completion due to MacNeille of the partially ordered space $(\mathcal{M}^0(\Omega), \leq)$ which was defined in (2.18), (2.19). Then this space $\mathcal{M}^0(\Omega)^\#$ in (2.20) is *order complete*, and we have the *order isomorphical embedding*, see Oberguggenberger & Rosinger [Appendix]

$$(2.21) \quad \mathcal{M}^0(\Omega) \ni G \longmapsto \langle G \rangle \in \mathcal{M}^0(\Omega)^\#$$

which also *preserves the infima and suprema*. Moreover, in view of the MacNeille construction, we can further extend the embedding (2.21) as follows

$$(2.22) \quad \begin{array}{ccccc} \mathcal{M}^0(\Omega) & \xrightarrow{\text{o. i. e.}} & \mathcal{M}^0(\Omega)^\# & \xrightarrow{\text{id}} & \mathcal{P}(\mathcal{M}^0(\Omega)) \\ G & \longrightarrow & < G] & \longrightarrow & < G] \end{array}$$

In order to obtain the full situation with respect to the range of the nonlinear partial differential operators $T(x, D)$, we note that $\mathcal{C}^0(\Omega) \subseteq \mathcal{C}_{nd}^0(\Omega)$, and we have the *order isomorphical embedding*

$$(2.23) \quad \mathcal{C}^0(\Omega) \ni g \longmapsto G \in \mathcal{M}^0(\Omega)$$

where G is the $\approx$ equivalence class of g . Furthermore, the partial order $\leq$ on $\mathcal{M}^0(\Omega)$ induces on $\mathcal{C}^0(\Omega)$ through this embedding the usual point-wise order relation of functions, namely, $g \leq h$, if and only if $g(x) \leq h(x)$, for $x \in \Omega$.

Let us now recall that our main interest is the construction of the commutative diagrams (1.9). In this regard, having constructed in (2.18) - (2.20), respectively, the spaces $\mathcal{M}^0(\Omega)$ and $\mathcal{M}^0(\Omega)^\#$ and their partial orders, the next step is to construct the partially ordered spaces $\mathcal{M}_T^m(\Omega)$ and $\mathcal{M}_T^m(\Omega)^\#$. For that purpose we start with (1.11), namely

$$(2.24) \quad T(x, D) : \mathcal{C}_{nd}^m(\Omega) \longrightarrow \mathcal{C}_{nd}^0(\Omega)$$

As mentioned, we shall solve the nonlinear PDEs in (1.1) by extending through order completion this mapping in (2.24), and we do so by constructing the commutative diagrams in (1.9). And at this stage we are now in the position to start doing so step by step. Let us note first that if $u \in \mathcal{C}^m(\Omega \setminus \Gamma)$, where $\Gamma \subset \Omega$ is any given closed, nowhere dense subset, then we also have

$$(2.25) \quad T(x, D)u \in \mathcal{C}^0(\Omega \setminus \Gamma)$$

This means that the singularity subsets Γ do *not* increase by the application of the nonlinear partial differential operators $T(x, D)$. However, as before with $\mathcal{C}_{nd}^0(\Omega)$, the ambiguity about associating such singularity sets to functions in $\mathcal{C}_{nd}^m(\Omega)$ remains. Therefore, in view of (2.25), we shall define the *equivalence* relation $u \approx_T v$, for any two functions $u, v \in \mathcal{C}_{nd}^m(\Omega)$, by the condition

$$(2.26) \quad T(x, D)u \approx T(x, D)v$$

which uses the equivalence relation $\approx$ given in (2.17), and in addition, it also depends on the nonlinear partial differential operator $T(x, D)$. In fact, this equivalence relation $\approx_T$ on $\mathcal{C}_{nd}^m(\Omega)$ is what is called the *pull-back* through the mapping $T(x, D)$ in (2.24) of the equivalence relation $\approx$ on $\mathcal{C}_{nd}^0(\Omega)$.

Let us now define the quotient space

$$(2.27) \quad \mathcal{M}_T^m(\Omega) = \mathcal{C}_{nd}^m(\Omega) / \approx_T$$

in which case the mapping (2.24) generates canonically the *injective* mapping

$$(2.28) \quad T : \mathcal{M}_T^m(\Omega) \longrightarrow \mathcal{M}^0(\Omega)$$

defined by $T(U) = G$, where G is the unique $\approx$ equivalence class in $\mathcal{M}^0(\Omega)$ of any of the $T(x, D)u$, where u belongs to the $\approx_T$ equivalence class U in $\mathcal{M}_T^m(\Omega)$.

At last, we can define the partial order $\leq_T$ on $\mathcal{M}_T^m(\Omega)$ as the *pull-back* through the mapping T in (2.28) of the partial order $\leq$ in (2.19) on $\mathcal{M}^0(\Omega)$, that is, for $U, V \in \mathcal{M}_T^m(\Omega)$, we have $U \leq_T V$, if and only if

$$(2.29) \quad T U \leq T V$$

In this way we obtain the partially ordered set $(\mathcal{M}_T^m(\Omega), \leq_T)$ giving the desired order structure on the domain $\mathcal{M}_T^m(\Omega)$ of T , which is the mapping in (2.28) that corresponds now to our nonlinear partial differential operator $T(x, D)$ in (1.2), (1.3), or more precisely, in (2.24). It is obvious in view of (2.29) that the injective mapping T in (2.28) is also an *order isomorphical embedding*.

So far, we have in this way obtained the top commutative rectangle in (1.9).

Applying now to $(\mathcal{M}_T^m(\Omega), \leq_T)$ the Dedekind order completion of Mac-Neille, we obtain

$$(2.30) \quad (\mathcal{M}_T^m(\Omega)^\#, \leq_T)$$

which is *order complete*, and in addition, similar with (2.21), we also have the *order isomorphical embedding*

$$(2.31) \quad \mathcal{M}_T^m(\Omega) \ni U \longmapsto \langle U \rangle \in \mathcal{M}_T^m(\Omega)^\#$$

which *preserves the infima and suprema*. Also, similar with (2.22), we have

$$(2.32) \quad \begin{array}{ccccc} \mathcal{M}_T^m(\Omega) & \xrightarrow{\text{o. i. e.}} & \mathcal{M}_T^m(\Omega)^\# & \xrightarrow{\text{id}} & \mathcal{P}(\mathcal{M}_T^m(\Omega)) \\ U & \longrightarrow & < U] & \longrightarrow & < U] \end{array}$$

And now all that remains is to define $T^\#$ in (1.9). In view of (2.32), however, this can be done in a standard manner following from the MacNeille order completion, see Oberguggenberger & Rosinger [Appendix]. Consequently, one obtains the *order isomorphical embedding*

$$(2.33) \quad T^\# : \mathcal{M}_T^m(\Omega)^\# \longrightarrow \mathcal{M}^0(\Omega)^\#$$

which also *preserves the infima and the suprema*. In more detail, we have the following commutative diagram

$$(2.34) \quad \begin{array}{ccc} \mathcal{M}_T^m(\Omega) \ni U & \xrightarrow{T} & T(U) \in \mathcal{M}^0(\Omega) \\ \downarrow & & \downarrow \\ \mathcal{M}_T^m(\Omega)^\# \ni < U] & \xrightarrow{T^\#} & T^\#(< U]) = \\ & & = < T(U)] \in \mathcal{M}^0(\Omega) \end{array}$$

In this way we have indeed obtained the whole of the commutative diagram (1.9), which we shall present now in the form seen next. Here "sur" and "inj" mean mappings which are surjective and injective, respectively, while as before, "o. i. e." means order isomorphical embedding, and "o. i." stands for order isomorphism. The dotted arrows " $< - - - - -$ " mean the "pull-back" through which the respective structures were defined

$$\begin{array}{ccc}
\mathcal{C}_{nd}^m(\Omega) & \xrightarrow{T(x, D)} & \mathcal{C}_{nd}^0(\Omega) \\
\downarrow \text{sur} \quad \approx_T & & \approx \quad \downarrow \text{sur} \\
\mathcal{M}_T^m(\Omega) & \xrightarrow[T]{\text{o. i. e.}} & \mathcal{M}^0(\Omega) \\
\downarrow \text{inj} \quad \leq_T & & \leq \quad \downarrow \text{inj} \\
\mathcal{M}_T^m(\Omega)^\# & \xrightarrow[T^\#]{\text{o. i., see (3.1) below}} & \mathcal{M}^0(\Omega)^\#
\end{array}
\tag{2.35}$$

3. General existence result

One of the typical *main existence results* concerning the solutions of the nonlinear PDEs in (1.1) is presented in the following theorem, see Oberguggenberger & Rosinger [38-64] for a proof

Theorem 3.1.

$$(3.1) \quad T^\# (\mathcal{M}_T^m(\Omega)^\#) = \mathcal{M}^0(\Omega)^\#$$

This means that, given the nonlinear PDEs in (1.1), for every right hand term $f \in \mathcal{M}^0(\Omega)^\#$, there exists a generalized solution $U \in \mathcal{M}_T^m(\Omega)^\#$, satisfying the relation $T^\# U = f$, according to the extension in (1.9).

As seen in Oberguggenberger & Rosinger [74-93], the space $\mathcal{M}^0(\Omega)^\#$ in which the right hand terms f of the nonlinear PDEs in (1.1) can range - and which now are solved by Theorem 3.1. - contains a large amount of *discontinuous* function on Ω . Certainly, in view of (2.18), $\mathcal{M}^0(\Omega)^\#$ contains all the *piecewise discontinuous* functions in $\mathcal{C}_{nd}^0(\Omega)$.

What is particularly important to note is that, in view of (3.1), a variety of linear and nonlinear PDEs can be solved, in spite of the fact that the respective PDEs are known *not* to have solutions in distributions. Among them is the celebrated 1957 Hans Lewy example, see Oberguggenberger & Rosinger [chap. 6, 8].

In this regard, it was in Oberguggenberger & Rosinger that this Hans Lewy example of a PDE not solvable in distribution was nevertheless solved for the first time through the method of order completion.

The *coherence* between the solutions obtained in (3.1) and the usual classical solutions, whenever the nonlinear PDEs in (1.1) may have the latter, follows easily from the commutative diagram (2.35). In other words, whenever the nonlinear PDEs in (1.1) happen to have classical solutions $U \in \mathcal{C}^m(\Omega)$, then they are also generalized solution in the sense of (3.1).

Finally, it is important to note that the above existence result in (3.1) can easily be extended to *systems* of nonlinear PDEs of the general form in (1.1), see Oberguggenberger & Rosinger [chap. 8-11].

4. Initial and/or boundary value problems and constitutive relations

One of the significant advantages of the order completion method in solving systems of nonlinear PDEs of the general form in (1.1) comes from the *ease* initial and/or boundary value problems associated with such equations can be solved. This is in strong contradistinction with the variety of functional analytic methods of solution where considerable difficulties arise related to the need to restrict distributions or generalized functions to lower dimensional manifolds. Indeed, such operations of restriction are typically ill-defined.

On the other hand, when using the method of order completion, the issue of satisfying the initial and/or boundary values can be *decoupled* from the issue of the existence of solutions. Indeed, satisfying the initial and/or boundary values can be dealt with *first* and *separately from* the issue of proving the existence of solutions.

Details in this regard can be found in Oberguggenberger & Rosinger

[chap. 8, 11]. And the reason behind that rather surprising ease the order completion method exhibits when dealing with initial and/or boundary value problems comes from a fact seen next, in section 5.

In Fluid Dynamics, and in general, Continuum Mechanics, a critical role is played by *constitutive equations*, see Rajagopal & Wineman, Rajagopal, or Rajagopal & Srinivasa.

The usual functional analytic methods can - if at all - deal with such constitutive equations in no less a difficult manner than they can do with initial and boundary value conditions. A regrettable consequence of these considerable difficulties is the failure so far of functional analytic methods to approach in any significant, let alone, systematic manner, the issue of such critically important constitutive equations. Here again, the order completion method proves its advantage by being able to deal as well with constitutive equations. This is simply a consequence of the fact that, as mentioned next in section 5, the order completion method can solve equations which are far more general than the linear or nonlinear systems of PDEs, or the constitutive equations.

5. An abstract existence result

A better understanding of the power underlying the order completion method in solving very large classes of equations, classes *far beyond* any nonlinear systems of PDEs, can be obtained from the following rather abstract existence result, see Oberguggenberger & Rosinger [chap. 9].

Let X be any set, and let $(Y, \leq)$ be any partially ordered set which has no minimum or maximum. Further, let

$$(5.1) \quad T : X \longrightarrow Y$$

be any given mapping. The problem we consider is to find a solution $A \in X$ for the equation

$$(5.2) \quad T(A) = F$$

for any given $F \in Y$. The answer is obtained as follows. Similar with the construction of the commutative diagrams (1.9) and (2.35), one can construct commutative diagrams

$$(5.3) \quad \begin{array}{ccc} X & \xrightarrow{T} & Y \\ \downarrow & & \downarrow \\ X_T & \xrightarrow{\tilde{T}} & Y \\ \downarrow & & \downarrow \\ X_T^\# & \xrightarrow[\text{bijective}]{T^\#} & Y^\# \end{array}$$

And then the following result on the existence of solutions holds

Theorem 5.1.

For any given $F \in Y^\#$, the equation

$$(5.4) \quad T^\#(A) = F$$

has a solution $A \in X_T^\#$, if and only if

$$(5.5) \quad \begin{aligned} \sup_{Y^\#} \{ T^\#(U) \mid U \in X_T^\#, \ T^\#(U) \subseteq F \} = \\ = \inf_{Y^\#} \{ T^\#(V) \mid V \in X_T^\#, \ F \subseteq T^\#(V) \} \end{aligned}$$

The significant generality of the above existence result allows, among others, the *separation* mentioned in section 4, between first satisfying the initial and/or boundary value conditions, and then, second, proving the existence of solutions in the case of general nonlinear systems of PDEs of the form in (1.1). Indeed, by first imposing the initial and/or boundary values, one is in fact defining the set X in (5.1). And as seen above, that can be done without any restrictions. Subsequently, condition (5.5) is both necessary and sufficient for the existence of a generalized solution $A \in X_T^\#$.

6. The Hausdorff continuity of solutions

The major *novelty* in this paper is about the fact that the solutions $U \in \mathcal{M}_T^m(\Omega)^\#$ of systems of nonlinear PDEs of type (1.1), obtained according to the procedure in Theorem 3.1., and of its generalizations can now be assimilated with *Hausdorff continuous functions* in $\mathbb{H}(\Omega)$. In fact, as seen in (A.12) in the Appendix, the mentioned solutions can be assimilated with *nearly finite Hausdorff continuous functions*.

7. Final comments

A further advantage of the order completion method is that one is *not* limited to consider in (1.9) and (2.35) only the *pull-back* partial order $\leq_T$ generated by the partial differential operators $T(x, D)$ on $\mathcal{M}_T^m(\Omega)$. Indeed, as seen in Oberguggenberger & Rosinger [chap. 13], a large variety of other partial orders on $\mathcal{M}_T^m(\Omega)$ can still secure existence theorems similar to Theorem 3.1.

As for the use of pull-back orders, it is important to note that there exists a certain analogy with functional analytic methods for solving PDEs. Indeed, in such methods, the topologies considered on the domains of the partial differential operators $T(x, D)$ are but pull-backs through these operators of suitable topologies on their ranges.

Details about such facts, and in general, about certain similarities between the order completion method and the usual functional analytic ones in solving PDEs can be found in Oberguggenberger & Rosinger [chap. 12].

The results in this paper invite a *comparison* with the *customary perception* regarding the solution of linear or nonlinear PDEs. Typical for that perception are the following to recent citations.

The 2004 edition of the Springer Universitext book "Lectures on PDEs" by V I Arnold, starts on page 1 with the statement :

"In contrast to ordinary differential equations, there is *no unified theory* of partial differential equations. Some equations have their own theories, while others have no theory

at all. The reason for this complexity is a more complicated geometry ...” (italics added)

Similarly, the 1998 edition of the book ”Partial Differential Equations” by L C Evans, starts his Examples on page 3 with the statement :

”There is no general theory known concerning the solvability of all partial differential equations. Such a theory is *extremely unlikely* to exist, given the rich variety of physical, geometric, and probabilistic phenomena which can be modelled by PDE. Instead, research focuses on various particular partial differential equations ...” (italics added)

Appendix : Definition and Properties of Hausdorff-Continuous Functions

The Hausdorff continuous functions are not unlike the usual real valued continuous functions. For instance, they assume real values on a dense subset of their domain of definition and are completely determined by the values on this subset. However, these functions may also assume *interval* values on a certain subset of their domain of definition. Hence the concept of Hausdorff continuity is formulated within the realm of *interval valued functions*. We shall deal in this Appendix with functions whose values can be not only usual real numbers but also *extended* real numbers, that is, elements in $\overline{\mathbb{R}} = \mathbb{R} \cup \{-\infty, +\infty\}$. Moreover, as mentioned it proves to be convenient to allow the values of the functions to be not only numbers in $\overline{\mathbb{R}}$, but also *closed intervals* of such numbers, namely, $[a, b] \subseteq \overline{\mathbb{R}}$, with $a, b \in \overline{\mathbb{R}}$, $a \leq b$.

Towards the end of the 19-th century, Baire brought in the concepts of *lower* and *upper semi-continuous* functions, when dealing with non-smooth real valued functions. And in effect, he associated with each real valued function f , *two* other real, or extended real valued functions $I(f)$ and $S(f)$, with $I(f) \leq f \leq S(f)$, which proved to be particularly helpful, see (A.6), (A.7). However, following the prevailing mentality at the time, each of these three functions was considered separately and as being a single valued function.

As it turns out on the other hand, by considering *interval valued* functions, such as for instance $F(f) = [I(f), S(f)]$, one can significantly improve on the understanding and handling of non-continuous functions.

The study of interval valued functions can, among others, show that the particular case of functions which have values given by one single number is appropriate for continuous functions only. On the other hand, non-continuous functions are much better described by suitably associated interval valued functions.

Indeed, in the case of functions f which are *not* continuous, a much better description can be obtained by considering them given by a *pair* of usual point valued functions, namely $f = [\underline{f}, \overline{f}]$, thus leading to interval valued functions, according to $f(x) = [\underline{f}(x), \overline{f}(x)] \subseteq \overline{\mathbb{R}}$, for $x \in \Omega$. And then, a natural class which replaces, and also extends, the usual point valued continuous functions is that of *Hausdorff-continuous* interval valued functions, see below Definition A1. The distinctive and *essential* feature of these Hausdorff-continuous functions $f = [\underline{f}, \overline{f}]$ is a condition of *minimality* with respect to the *gap* between $\underline{f}$ and $\overline{f}$, with the further requirement that $\underline{f}$ be lower semi-continuous, and $\overline{f}$ be upper semi-continuous.

The interest in more recent times in interval valued functions comes from a number of branches of mathematics, such as approximation theory, Sendov, and numerical analysis, Kraemer.

Most of the results on interval valued functions presented in this Appendix have, however, been developed by R Anguelov. This was done in view of the usefulness of such functions in several branches of mathematics, see Anguelov [1,2], Anguelov & Markov, Anguelov et.al [1,2], Anguelov & Minani, Anguelov & Rosinger [1-3]. Here, owing to restriction of space, we shall only present a minimal amount of them, needed in order to support section 6 above. For the full details in this regard, including proofs, see Appendix 2 in Anguelov & Rosinger [3].

As mentioned, the class of interval valued functions of special interest here is that of *Hausdorff-continuous*, or in short, *H-continuous* functions. As it turns out they enjoy the *minimality* property (A.8) with respect to their *graph completion*, see (A.7), and that allows an effective interplay between Analysis and Topology, with the latter in-

volving both the domain and the range of the functions dealt with. Let

$$(A.1) \quad \overline{\mathbb{IR}} = \{ [\underline{a}, \bar{a}] \mid \underline{a}, \bar{a} \in \overline{\mathbb{R}} = \mathbb{R} \cup \{-\infty, +\infty\}, \underline{a} \leq \bar{a} \}$$

be the set of all finite or infinite closed intervals. The functions which we consider can be defined on arbitrary topological spaces Ω . For the purposes of the nonlinear PDEs studied in this paper, however, it will be sufficient to assume that $\Omega \subseteq \mathbb{R}^n$ are arbitrary open subsets. Let us now consider the set of interval valued functions

$$(A.2) \quad \mathbb{A}(\Omega) = \{ f : \Omega \longrightarrow \overline{\mathbb{IR}} \}$$

By identifying the point $a \in \overline{\mathbb{R}}$ with the degenerate interval $[a, a] \in \overline{\mathbb{IR}}$, we consider $\overline{\mathbb{R}}$ as a subset of $\overline{\mathbb{IR}}$. In this way $\mathbb{A}(\Omega)$ will contain the set of functions with extended real values, namely

$$(A.3) \quad \mathcal{A}(\Omega) = \{ f : \Omega \longrightarrow \overline{\mathbb{R}} \} \subseteq \mathbb{A}(\Omega)$$

We define a partial order $\leq$ on $\overline{\mathbb{IR}}$ by

$$(A.4) \quad [\underline{a}, \bar{a}] \leq [\underline{b}, \bar{b}] \iff \underline{a} \leq \underline{b}, \bar{a} \leq \bar{b}$$

Now on $\mathbb{A}(\Omega)$ we define the partial order induced by (A.4) in the usual point-wise way, namely, for $f, g \in \mathbb{A}(\Omega)$, we have

$$(A.5) \quad f \leq g \iff f(x) \leq g(x), \quad x \in \Omega$$

Clearly, when restricted to $\mathcal{A}(\Omega)$, the above partial order on $\mathbb{A}(\Omega)$ reduces to the usual one among point valued functions.

Let $f \in \mathbb{A}(\Omega)$. For every $x \in \Omega$, the value of f is an interval, namely, $f(x) = [\underline{f}(x), \bar{f}(x)]$, with $\underline{f}(x), \bar{f}(x) \in \overline{\mathbb{R}}$, $\underline{f}(x) \leq \bar{f}(x)$. Hence, every function $f \in \mathbb{A}(\Omega)$ can be written in the form $f = [\underline{f}, \bar{f}]$, with $\underline{f}, \bar{f} \in \mathcal{A}(\Omega)$, $\underline{f} \leq f \leq \bar{f}$, and $f \in \mathcal{A}(\Omega) \iff \underline{f} = f = \bar{f}$.

In the particular case of functions in $\mathcal{A}(\Omega)$, that is, with extended real, but point, and not nondegenerate interval values, a number of basic results were obtained already in Baire, see also Nicolescu for a more recent detailed presentation. The rest of the more general results concerning functions in $\mathbb{A}(\Omega)$, that is, with values finite or infinite closed intervals, were developed for the first time in the above cited works

of Anguelov . The few such earlier results were obtained in Sendov, where the particular instance of $\Omega \subseteq \mathbb{R}$ was dealt with.

For $x \in \Omega$, we denote by $B_\delta(x)$ the open ball of radius δ centered at x . Let us consider any *dense* subset $D \subseteq \Omega$, and associate with it the pair of mappings $I(D, \Omega, \cdot), S(D, \Omega, \cdot) : \mathbb{A}(\Omega) \rightarrow \mathcal{A}(\Omega)$, called *lower* and *upper Baire operators*, respectively, where for every function $f \in \mathbb{A}(\Omega)$ and $x \in \Omega$, we define

$$(A.6) \quad \begin{aligned} I(D, \Omega, f)(x) &= \sup_{\delta > 0} \inf \{ z \in f(y) \mid y \in B_\delta(x) \cap D \} \\ S(D, \Omega, f)(x) &= \inf_{\delta > 0} \sup \{ z \in f(y) \mid y \in B_\delta(x) \cap D \} \end{aligned}$$

In Baire, these two operators were considered and studied in the particular case of functions $f \in \mathcal{A}(\Omega)$ and when $D = \Omega$, see also Nicolescu. In view of the main interest here in *interval valued* functions $f \in \mathbb{A}(\Omega)$, it is useful to consider as well the following third mapping, namely, $F(D, \Omega, f) : \mathbb{A}(\Omega) \rightarrow \mathbb{A}(\Omega)$, defined for $f \in \mathbb{A}(\Omega)$ by

$$(A.7) \quad F(D, \Omega, f)(x) = [I(D, \Omega, f)(x), S(D, \Omega, f)(x)], \quad x \in \Omega,$$

and called the *graph completion operator*. In case $D = \Omega$, we use the simpler notations $I(\Omega, \Omega, f) = I(f)$, $S(\Omega, \Omega, f) = S(f)$ and $F(\Omega, \Omega, f) = F(f)$.

The next definition was given in Sendov in the case of $\Omega \subseteq \mathbb{R}$, however, it can obviously be extended to any topological space Ω .

Definition A.1

A function $f \in \mathbb{A}(\Omega)$ is called *Hausdorff-continuous*, or in short, *H-continuous*, if and only if for every function $g \in \mathbb{A}(\Omega)$, we have satisfied the following *minimality* condition on f

$$(A.8) \quad g(x) \subseteq f(x), \quad x \in \Omega \implies F(g) = f$$

We shall denote by $\mathbb{H}(\Omega)$ the set of all Hausdorff-continuous interval valued functions on Ω .

Definition A2

A function $f \in \mathbb{H}(\Omega)$ is called *nearly finite*, if and only if there exists an open and dense subset $D \subseteq \Omega$, such that

$$(A.9) \quad f(x) \in \overline{\mathbb{I}\mathbb{R}} \quad \text{is a finite interval for } x \in D$$

We denote by $\mathbb{H}_{nf}(\Omega)$ the set of nearly finite H-continuous functions $f \in \mathbb{A}(\Omega)$.

Regarding the *regularity* properties of solutions of general nonlinear systems of PDEs of the form in (1.1), a crucial role is played by the following mapping

$$(A.10) \quad F_0 : \mathcal{C}_{nd}^0(\Omega) \ni u \longmapsto F(\Omega \setminus \Gamma, \Omega, u) \in \mathbb{H}_{nf}(\Omega)$$

where we recall that, according to (1.7), for every $u \in \mathcal{C}_{nd}^0(\Omega)$, there exists a closed, nowhere dense subset $\Gamma \subset \Omega$, such that $u \in \mathcal{C}^0(\Omega \setminus \Gamma)$, hence in view of (A.7), $F(\Omega \setminus \Gamma, \Omega, u)$ is well defined. The fact that such a Γ need not be unique, does not affect the above definition, see Anguelov & Rosinger [3].

The following theorem shows that the images of two functions in $\mathcal{C}_{nd}(\Omega)$ under the mapping F_0 in (A.10) are the same, if and only if these functions are equivalent with respect to the equivalence relation (2.17).

Theorem A.1

Let $u, v \in \mathcal{C}_{nd}(\Omega)$. Then $F_0(u) = F_0(v) \iff u \approx v$

In view of (2.18), (A.10) and the above theorem now we can define a mapping

$$(A.11) \quad \tilde{F}_0 : \mathcal{M}^0(\Omega) \longrightarrow \mathbb{H}_{nf}(\Omega)$$

in the following way. Let $u \in U \in \mathcal{M}^0(\Omega)$, then $\tilde{F}_0(U) = F(u)$. It is easy to see that the definition of $\tilde{F}_0(U)$ does not depend on the particular representative $u \in U$ of the equivalence class U .

Theorem A.2

The mapping $\tilde{F}_0 : \mathcal{M}^0(\Omega) \longrightarrow \mathbb{H}_{nf}(\Omega)$ defined in (A.11) is an *order isomorphic embedding* with respect to the order relation (2.19) on $\mathcal{M}^0(\Omega)$ and the order relation induced by (A.5) on $\mathbb{H}_{nf}(\Omega)$. Namely, for any $U, V \in \mathcal{M}^0(\Omega)$, we have

$$U \leq V \iff \tilde{F}_0(U) \leq \tilde{F}_0(V)$$

Finally, we also have

Theorem A.3

The set $\mathbb{H}_{nf}(\Omega)$ is Dedekind order complete with respect to the partial order induced on it by (A.5).

Let $g \in \mathbb{H}_{nf}(\Omega)$. Then there exists a subset $\mathcal{G} \subseteq \mathcal{M}^0(\Omega)$ such that

$$g = \sup \tilde{F}_0(\mathcal{G}) = \sup \{ \tilde{F}_0(G) \mid G \in \mathcal{G} \}$$

This theorem shows that $\mathbb{H}_{nf}(\Omega)$ is the *smallest* Dedekind order complete subset of $\mathbb{H}(\Omega)$ which contains the image of $\mathcal{M}^0(\Omega)$ under the order isomorphical embedding $\tilde{F}_0$. Hence it is *order isomorphic* to the Dedekind order completion $\mathcal{M}^0(\Omega)^\#$ of $\mathcal{M}^0(\Omega)$. In this way we obtain the commutative diagram, where $\tilde{F}_0^\#$ denotes the order isomorphism from $\mathcal{M}^0(\Omega)^\#$ to $\mathbb{H}_{nf}(\Omega)$, namely

$$\begin{array}{ccc}
 C_{nd}^0(\Omega) & \xrightarrow[\text{graph completion}]{F_0} & \mathbb{H}_{nf}(\Omega) \\
 \downarrow & & \updownarrow \\
 \mathcal{M}^0(\Omega) = C_{nd}^0(\Omega)/\approx & \xrightarrow[\text{order isomorphical embedding}]{\tilde{F}_0} & \mathbb{H}_{nf}(\Omega) \\
 \downarrow & & \updownarrow \\
 \mathcal{M}^0(\Omega)^\# & \xrightarrow[\text{order isomorphism}]{\tilde{F}_0^\#} & \mathbb{H}_{nf}(\Omega)
 \end{array}$$

Now we can bring together the above diagram with the one in (2.35) and obtain the two successive *order isomorphisms*

$$(A.12) \quad \mathcal{M}_T^m(\Omega)_T^\# \xrightarrow{T^\#} \mathcal{M}^0(\Omega)^\# \xrightarrow{\tilde{F}_0^\#} \mathbb{H}_{nf}(\Omega)$$

It follows therefore that the set $\mathcal{M}_T^m(\Omega)_T^\#$ in which the *solutions* of the general nonlinear systems of PDEs of the form in (1.1) are found is mapped by the *bijection* $\tilde{F}_0^\# \circ T^\#$ onto the set $\mathbb{H}_{nf}(\Omega)$ of all *nearly finite Hausdorff continuous functions*. Since both these mappings are order isomorphisms, the set $\mathcal{M}_T^m(\Omega)_T^\#$ is order isomorphic with the set $\mathbb{H}_{nf}(\Omega)$.

Hence, the solutions of the general nonlinear systems of PDEs of

the form in (1.1), which are obtained through the order completion method, can always be assimilated with nearly finite Hausdorff continuous functions.

This is the argument supporting section 6 above.

For the sake of further clarification, let us turn now in the remaining part of this Appendix to some of the issues concerning the *discontinuities* of Hausdorff-continuous functions. Arbitrary interval valued functions $f = [\underline{f}, \overline{f}] \in \mathbb{A}(\Omega)$ can exhibit a variety of types of discontinuities, and certainly not less so, than usual point valued functions $f \in \mathcal{A}(\Omega)$ do.

Hausdorff-continuous functions, although generalize usual point valued continuous functions, enjoy nevertheless a number of nontrivial *continuity* related properties. On the other hand, Hausdorff-continuous functions can have quite large sets of *discontinuities*. This shows that they do indeed form a larger class than the usual continuous functions, even if they still have important similar properties.

The fact that large enough sets of discontinuities can be present with Hausdorff-continuous functions allows for their use - as seen in this paper - in obtaining the *existence of nonclassical solutions* for large classes of systems of nonlinear PDEs.

Consequently, and as mentioned, Hausdorff-continuous functions - precisely since they are *not* generalized functions - can be seen as *setting aside* to a certain extent the variety of distributional and other traditional generalized solutions of linear and nonlinear PDEs which have been obtained by functional analytic methods, or by the methods of the nonlinear algebraic theory listed by the AMS Subject Classification 2000, under 46F30.

Indeed, solving large classes of nonlinear PDEs through Hausdorff-continuous functions offers, among others, the following *double advantage* :

- one can bring in a significant *simplification* by avoiding the variety of usual functional analytic methods with their spaces of distributions or generalized functions, and
- one can obtain *universal regularity* results for solutions of large classes of systems of nonlinear PDEs.

Theorems A.4 and A.5 below show important properties of *Hausdorff-continuous* functions related to their sets of *discontinuities*. For every interval valued function $f = [\underline{f}, \overline{f}] \in \mathbb{A}(\Omega)$, we denote by $\Gamma(f) = \{ x \in \Omega \mid \underline{f}(x) < \overline{f}(x) \}$, which is the set of points $x \in \Omega$ where f assumes values $f(x) = [\underline{f}(x), \overline{f}(x)]$ that are *non-degenerate intervals*, and not merely points.

It follows that at points $x \in \Gamma(f)$, the interval valued function $f = [\underline{f}, \overline{f}]$ *cannot* be continuous in the usual sense, since it is not a usual point valued function.

In the particular case of *Hausdorff-continuous* functions, this fact can further be clarified. Namely, given any point $x \in \Omega$, then

$$(A.13) \quad \begin{aligned} x \in \Gamma(f) &\iff \underline{f} \text{ and } \overline{f} \text{ not continuous at } x \iff \\ &\iff \underline{f} \text{ or } \overline{f} \text{ not continuous at } x \iff \underline{f}(x) < \overline{f}(x) \end{aligned}$$

And now the basic result on the discontinuities of Hausdorff-continuous functions

Theorem A.4

Given any H-continuous function $f \in \mathbb{A}(\Omega)$. Then $\Gamma(f)$ is of *first Baire category* in Ω .

Let us further specify the structure of the discontinuity set $\Gamma(f)$. For $\epsilon > 0$, let us denote $\Gamma_\epsilon(f) = \{ x \in \Omega \mid \overline{f}(x) - \underline{f}(x) \geq \epsilon \}$. Then clearly $\Gamma(f) = \bigcup_{\epsilon > 0} \Gamma_\epsilon(f) = \bigcup_{n \geq 1} \Gamma_{1/n}(f)$.

The next theorem gives a further insight into the structure of the discontinuity set $\Gamma(f)$ of Hausdorff-continuous functions $f \in \mathbb{A}(\Omega)$.

Theorem A.5

If the function $f \in \mathbb{A}(\Omega)$ is H-continuous, then for every $\epsilon > 0$, the set $\Gamma_\epsilon(f)$ is *closed* and *nowhere dense* in Ω .

An important *similarity* between usual continuous, and on the other hand, Hausdorff-continuous functions is that both of them are determined *uniquely* if they are known on a *dense* subset of their domains of definition. This property comes in spite of the fact that, as seen in the previous section, Hausdorff-continuous functions can have discontinuities on sets of first Baire category, and such sets can have arbitrary large positive Lebesgue measure, see Oxtoby. Indeed, we have

Theorem A.6

Let $f = [\underline{f}, \overline{f}]$, $g = [\underline{g}, \overline{g}] \in \mathbb{A}(\Omega)$ be two H-continuous functions, and suppose given any dense subset $D \subseteq \Omega$. Then

$$f(x) = g(x), \quad x \in D \implies f = g \text{ on } \Omega$$

The real line $\mathbb{R}$ is Dedekind order complete, but not order complete, while the extended real line $\overline{\mathbb{R}}$ is both Dedekind order complete and order complete.

Let us recall that a partially ordered set which is order complete will also be Dedekind order complete, but as seen above, not necessarily the other way round as well.

Typically, various spaces of real valued functions encountered in Analysis are neither Dedekind order complete, nor order complete, when considered with the natural point-wise partial order relation.

However, as we can see next, this situation changes when we deal with the set $\mathbb{H}(\Omega)$ of Hausdorff-continuous functions.

Theorem A.7 (Anguelov [2])

The set $\mathbb{H}(\Omega)$ of Hausdorff-continuous functions is order complete when considered with the partial order in (A.5).

The Dedekind order completeness of the space $\mathbb{H}(\Omega)$ of Hausdorff-continuous functions is a *nontrivial* property, in view of the various connections between the usual continuous, and on the other hand Hausdorff-continuous functions. Indeed, the space $\mathcal{C}(\Omega)$ of usual real valued continuous functions, which we have seen is strictly contained in $\mathbb{H}(\Omega)$, is well known *not* to be Dedekind order complete. On the other hand, once $\mathbb{H}(\Omega)$ proves to be Dedekind order complete, its order completeness follows easily from the fact that $\overline{\mathbb{R}}$ is order complete. Indeed, the smallest and largest elements in $\mathbb{H}(\Omega)$ are respectively the functions $\Omega \ni x \mapsto -\infty$ and $\Omega \ni x \mapsto +\infty$.

As is well known and shown by simple examples the spaces of real valued continuous functions $\mathcal{C}(\Omega)$ are *not* Dedekind order complete, thus, are not order complete either.

Since these spaces are partially ordered in a natural way, one can apply to them the MacNeille version of Dedekind order completion method, see Oberguggenberger & Rosinger [Appendix], Luxemburg & Zaanen,

or Zaanen.

This however being a general construction based on Dedekind type cuts, it leaves open the question of the *nature* of the elements which are added to these spaces of continuous functions by the respective Dedekind order completion process.

A classical, 1950 result in this regard was obtained by Dilworth in the case of *bounded* and real valued continuous functions on Ω for arbitrary completely regular topological spaces Ω . Namely, the respective Dedekind order completion is given by all the normal upper semi-continuous functions on Ω . Regarding the Dedekind order completion of spaces $\mathcal{C}(\Omega)$ certain results were obtained in Mack & Johnson.

As seen in Anguelov [1], the Dedekind order completion of spaces $\mathcal{C}(\Omega)$ of real valued continuous functions was for the first time effectively constructed for a large class of topological spaces Ω . In this construction Hausdorff-continuous functions and some of their subspaces play a crucial role. Consequently, both the problem of the *completion* of the spaces $\mathcal{C}(\Omega)$, as well as that of the *structure* of the elements which are added to these spaces $\mathcal{C}(\Omega)$ through the respective completion find a convenient solution through the use of *interval valued* functions.

References

- [1] Anguelov R [1] : Dedekind order completion of $C(X)$ by Hausdorff continuous functions. Quaestiones Mathematicae, Vol. 27, 2004, 153-170
- [2] Anguelov R [2] : An introduction to some spaces of interval functions. arXiv:math.GM/0408013
- [3] Anguelov R, Markov S : Extended segment analysis. Freiburger Intervall - Berichte 10, 1981, 1 - 63.
- [4] Anguelov R, Markov S, Sendov B [1] : On the Normed Linear Space of Hausdorff Continuous Functions. Poceedings of the Fifth International Conference on "Large Scale Scientific Computations", June 6-10, 2005, Sozopol, Lecture Notes in Computer Science, Springer (to appear)

- [5] Anguelov R, Markov S, Sendov B [2] : The Linear Space of Hausdorff Continuous Functions, Technical Report UPWT2004/4
- [6] Anguelov R, Minani F : Interval Viscosity Solutions of Hamilton-Jacobi Equations. Technical Report UPWT 2005/3, University of Pretoria
- [7] Anguelov R, Rosinger E E [1] : Dedekind order completion of $\mathcal{M}(\Omega)$ by Hausdorff continuous functions. (to appear)
- [8] Anguelov R, Rosinger E E [2] : Hausdorff Continuous Solutions of Nonlinear PDEs through the Order Completion Method. *Quaestiones Mathematicae* (to appear), see arXiv:math.AP/0406517
- [9] Anguelov R, Rosinger E E [3] : Solution of Nonlinear PDEs by Hausdorff Continuous Functions (to appear).
- [10] Arnold V I : Lectures on PDEs. Springer Universitext, 2004
- [11] Baire R : *Lecons sur les Fonctions Discontinues*. Collection Borel, Paris, 1905.
- [12] Bardi M, Capuzzo-Dolcetta I : Optimal control and viscosity solutions of Hamilton-Jacobi-Bellman equations. Birkhäuser, Boston, Basel, Berlin, 1997.
- [13] Birkhoff G : The Role of Order in Computing. In Moore R (Ed.) : *Reliability in Computing*. Academic Press, 1988, 357–378.
- [14] Dilworth R P : The normal completion of the lattice of continuous functions. *Trans. Amer. Math. Soc.*, 68, 1950, 427–438.
- [15] Evans L C : *Partial Differential Equations*. AMS Graduate Studies in Mathematics, Vol. 19, 1998
- [16] Forster O : *Analysis 3, Integralrechnung in $\mathbb{R}^n$ mit Anwendungen*. Friedr. Vieweg, Braunschweig, Wiesbaden, 1981
- [17] Kraemer W, von Gudenberg J W (Eds) : *Scientific Computing, Validated Numerics, Interval Methods*. Kluwer, Dordrecht, 2001
- [18] Lewy, H : An example of smooth linear partial differential equation without solutions. *Ann. Math.*, vol. 66, no. 2, 1957, 155-158

- [19] Luxemburg W A J, Zaanen A C : Riesz Spaces I. North Holland, Amsterdam, 1971.
- [20] Mack J E, Johnson D G : The Dedekind completion of $C(X)$. Pacif. J. Math., 20, 2, 1967, 231-243
- [21] MacNeille H M : Partially ordered sets. Trans. AMS, vol. 42, 1937, 416-460
- [22] Markov S : A nonstandard subtraction of intervals. Serdica. Vol. 3, 1977, 359-370
- [23] Markov S : Calculus for interval functions of a real variable. Computing, Vol. 22, 1979, 325-337
- [24] Markov S : Extended interval arithmetic involving infinite intervals. Mathematica Balkanica, Vol.6, 1992, 269-304.
- [25] Nicolescu M : Analiză Matematică II. Editura Technică, București, 1958
- [26] Oberguggenberger M B, Rosinger E E : Solution of Continuous Nonlinear PDEs through Order Completion. North-Holland Mathematics Studies, Vol. 181. North-Holland, Amsterdam, 1994
see also review MR 95k:35002
- [27] Oxtoby J C : Measure and Category. Springer, New York, 1971
- [28] Rajagopal K R, Wineman A S : On constitutive equations for branching of response with selectivity. International Journal of Nonlinear Mechanics, Vol. 15, 1980, 83-91
- [29] Rajagopal K R : On implicit constitutive theories. Application of Mathematics, Vol. 28, No. 4, 2003, 279-319
- [30] Rajagopal K R, Srinivasa A R : On thermo-mechanical restrictions of continua. Proc. R. Soc. London A Math., Vol. 460, 2004, 631-651
- [31] Rosinger E E : Hausdorff continuous solutions of arbitrary continuous nonlinear PDEs through the order completion method. arXiv:math.AP/0405546

- [32] Rosinger E E : Can there be a general nonlinear PDE theory for the existence of solutions ? arXiv:math.AP/0407026
- [33] Rosinger E E, Rudolph M : Group invariance of global generalised solutions of nonlinear PDEs : A Dedekind order completion method. Lie Groups and their Applications, Vol. 1, No. 1, July-August 1994, 203-215
- [34] Sendov B : Hausdorff Approximations. Kluwer, Dordrecht, 1990
- [35] Tutschke W : Initial Value Problems in Classes of Generalized Analytic Functions. Springer, New York, 1989
- [36] Zaanen A C : The universal completion of an Archimedean Riesz space. Indag. Math., 45, 4, 1983, 435-441
- [37] Zaharov V : Functional characterization of absolute and Dedekind completion. Bull. Acad. Polon. Sci., 29, 5-6, 1981, 293-297